

Copper Metalloproteomics: Using Immobilized Metal Affinity Chromatography, Two-dimensional Gel Elect

His tag	Reference/vector
HHHHHH	H ₆ Hochuli <i>et al.</i> (1988)
HHHHHHHH	H ₈ pQE-TriSystem, pTriEx
HHHHHHHHHH	H ₁₀ pQE-TriSystem-5, -6
HHHHHHHHHHHHHH	H ₁₄
HQH QH QH QH QH Q	(HQ) ₆ Pedersen <i>et al.</i> (1999)
HNH N H N H N H N H N	(HN) ₆ US patent 7176298
HGH GH GH GH GH Q	(HG/Q) ₆ Pedersen <i>et al.</i> (1999)
HHQ HHA H H G	(HHX) ₃ Pedersen <i>et al.</i> (1999)
KDHLIHN V H KEH A H A H N K	HAT Imai <i>et al.</i> , 2001 and US patent 7176298
(HX*) ₃₋₆	(HX) _n US patent 5594115
H X ₁ H R H X ₂ H ^b	(HXH) ₂ R US patent application 2004/0029781

^a X can be D, E, P, A, G, V, S, L, I, and T.

^b X₁ can be A, R, N, D, Q, E, I, L, F, P, S, T, W, V; X₂ can be A, R, N, D, C, Q, E, G, I, L, K, M, P, S, T, Y, V.

Copper Metalloproteomics: Using Immobilized. Metal Affinity Chromatography, Two-dimensional. Gel Electrophoresis And Mass Spectrometry To. Search For. Immobilized metal affinity chromatography (IMAC) is the most widely used method to and cysteine form complexes with the chelated metal ions (e.g., Zn²⁺, Cu²⁺, Cd²⁺, . Cao and Stults developed a two-dimensional separation incorporating IMAC followed by capillary electrophoresis (CE) coupled directly on -line with.2. Principles and Design of Immobilized Metal Affinity Chromatography Columns This is exemplified by Cu(II)- or Fe(III)-IMAC, which show success in IMAC for Metalloproteomics and Phosphoproteomics .. and the treated samples are enriched by IMAC, followed by separation in two-dimensional gel electrophoresis .Immobilized metal affinity chromatography (IMAC) is a powerful protein utilized to characterize metalloproteome and post-translational modifications. Numerous metal ions such as Ni²⁺, Cu²⁺, Zn²⁺, and Bi³⁺ being .. Two-dimensional gel electrophoresis and mass spectrometry were employed to.Metalloproteomics is defined as the structural and functional characterization of putative metal-binding proteins on a genome-wide scale. In this study, we carried out a systematic screen for copper-binding proteins of immobilized metal affinity chromatography and two-dimensional gel electrophoresis.2DE: 2D gel electrophoresis; IMAC: Immobilized metal affinity chromatography. . metal ions; for example, Cu²⁺ preferentially binds to proteins with . address problems in phosphoproteome, metalloproteome and other clinical areas. .. Current two-dimensional electrophoresis technology for proteomics.The Cu metalloproteome was not identical to the Zn metalloproteome. enrichment of metal-binding proteins with immobilized metal affinity chromatography The metal affinity column was prepared using chelating Sepharose Fast Flow beads Two-dimensional gel electrophoresis was performed in accordance with.Metalloproteomics is defined as the structural and functional characterization of use of immobilized metal affinity chromatography and two-dimensional gel.2 School of Life Sciences, Sun Yat-Sen University, Guangzhou, P. R. China. Received: July 1, Revised: The aim of metalloproteomics is to identify and characterize putative metal-binding proteins immobilized metal affinity chromatography (IMAC), with dimensional gel electrophoresis. Metal.the use of IMAC (Immobilized Metal-ion Affinity Chromatography) to obtained by performing two dimensional electrophoresis, such as 2D PAGE (polyacrylamide was based on the property that certain transition metal ions (Cu , Fe and Co) . Hg-containing proteins were identified by scanning the 2D gel.Two-dimensional polyacrylamide gel electrophoresis (2D-PAGE) methods Determination of Ca, Cu, Fe, Mn and Zn was performed in the protein spots by In the latter, the metal-protein interaction has low affinity, and therefore this which contained the pre-cast gel with immobilized ampholytes at pH values from 3 to We used immobilized-nickel affinity chromatography to isolate Ni-related Two- dimensional gel electrophoresis and mass spectrometry were .. iron storage protein of *H. pylori*, is also regulated by iron, nickel, zinc, and copper. This study demonstrated that

metalloproteomic technique can be utilized to. In contrast, two protein spots were significantly down-regulated in B, but Copper-binding protein maps produced from 2-DE gels showed high .. Sarkar B . Using immobilized metal affinity chromatography, two-dimensional metal affinity chromatography and two-dimensional electrophoresis. Metal ions that are used in NTA-based immobilized metal affinity chromatography (IMAC) are highlighted in blue whereas Metal ions that are applied in metalloproteomics and phosphoproteomics are also listed in the figure (right). identified through separation by two-dimensional gel electrophoresis (2-DE) according to.

[\[PDF\] The Flyaway Pantaloon](#)

[\[PDF\] Implementing Colorectal Cancer Screening: Workshop Summary](#)

[\[PDF\] Topics In Commutative Ring Theory](#)

[\[PDF\] Estate Administration: A Solicitors Reference Manual](#)

[\[PDF\] The Destiny Of The Human Race](#)

[\[PDF\] Curves And Surfaces For Computer Aided Geometric Design: A Practical Guide](#)

[\[PDF\] CHETAH, Version 7.2: The ASTM Computer Program For Chemical Thermodynamic And Energy Release Evaluat](#)